

Product Information

ASCOMILK F

Milkreplacer for calves

Description:

ASCOMILK F is not just a simple milk substitute. It's a special formula composed of dairy products, easily digestible fat, as well as carefully selected vitamins and minerals to promote growth of calves.

ASCOMILK F contains a small quantity of skim milk powder which gives it high acceptance.

ASCOMILK F is the key for a growth program for future highly productive dairy cows, as well as an efficient way to reduce rising cost.

Composition:

Whey powder, Vegetable oil, Soya flour, Skim milk powder, Wheat gluten, Lactose, Maltodextrin, Hydrolysed yeast, Wheat starch, Dicalcium phosphate, Magnesium sulphate, Magnesium hydroxide, Carbonate of calcium.

Advantages:

Milk protein	High digestibility: eases digestion in the first phase. Excellent performance.
Specially selected fat	Brings energy for a quick growth.
Vegetable protein	Economic rearing. Absorbable vegetable protein leads to a quick move to solid feed.
Vitamins and minerals	For an optimal growth and a complete nutrition.
Digestive safety	Through a supply of probiotics (<i>Enterococcus Faecium</i>), which acts as intestinal flora barrier. High level of protection from diarrhea.
Economic formula	Reduces production costs.

Typical analysis:

Moisture	5,0 %
Crude protein	21,0 %
Crude fat	16,0 %
Ashes	7,0 %
Crude fibre	0,9 %
Calcium	0,8 %
Sodium	0,5 %
Phosphor	0,7 %

Added vitamins and trace elements per kg:

Vitamin A	25.000 IU	Iron	40 mg
Vitamin D3	10.000 IU	Copper	12 mg
Vitamin E	80 IU	Zinc	70 mg
Vitamin C	180 IU	Manganese	65 mg
Vitamin K3	5 mg	Iode	0,3 mg
Vitamin B1	15 mg	Selenium	0,2 mg

Gut flora stabiliser

E1708 *Enterococcus Faecium* NCIMB 11181 1,3*10⁹ CFU/kg

Recommended nursing plan for rearing calves

2 meals / day		
Age / days	water	milk
1-4 days	COLOSTRUM	
5-7 days	2 l	240 g
week 2	2,5 l	350 g
week 3	3,0 l	420 g
week 4	3,0 l	420 g
week 5	3,0 l	420 g
week 6	3,0 l	420 g
week 7	3,0 l	420 g
week 8	2,0 l	280 g
week 9	1,2 l	210 g
week 10	WEANING	

After the second week: use straw or hay and water without limit with ASCOMILK F.

Preparation:

Mix ASCOMILK F with half the volume of hot water at 50°C and complete with warm water – feed at 38-40° C.

Use a tool to homogenize thoroughly and get a perfect solubility.

If you use a blending machine, do not use it for an extended period of time.

Set the thermostat to 40° C.

Advice:

- Do not make the preparation directly in the teat bucket. It is recommended to do the preparation aside and then pour it into the teat bucket.
- Only use potable water.
- Hygiene is important – clean all utensils after each feeding.
- To keep the milk replacer powder fresh, tightly roll the bag closed and store it in a dry and cool place.
- The feeding time schedule of the milk replacer must be fixed, with regular intervals.